Read enclosed leaflet before use. प्रयोग से पहले संलग्न लीफलेट पढ़ें।

SANTIVO - Tebuconazole 50% + Trifloxystrobin 25% WG (Systemic Fungicide)

Chemical Composition : 1 Tebuconazole a.i 50.00 % w/w, 2 Trifloxystrobin a.i. 25.00 % w/w, 3 Formaldehyde, reaction products with sulfonated 1,1- oxybis (methyl benzene) sodium salt 15.00 % w/w, 4 Naphthalenesulfonic acids, branched and linear butyl derivatives, sodium salt : 1.00 % w/w, 5 Ureaformaldehydecondensate 3.00 % w/w, 6 Kaolin Q.S. %, Total 100 % w/w Direction Of Use : Please read enclosed leaflet before use. Precautions : Protective clothing should be worn while mixing and application. Wash contaminated clothes and body part after application. avoid inhalation do not smoke, drink, eat &chew anything during application. **Antidote** : No specific antidote is known. Give symptomatic treatment. **First Aid** : 1. If ingested: If person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious person. Call the doctor. Do not induce vomiting until and unless desired by attending physician. 2. If inhaled: Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, preferably with mouth-to-mouth breathing. Consult a physician. 3. In case of skin contact: Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All the contaminated clothes should be segregated and thoroughly washed with soap and water separately before use. 4. In case of eye contact: Rinse gently with plenty of water for at least 15 minutes. If pain, irritation, redness or photophobia persists, consult eye specialist. Symptoms : Weakness, fatigue, despoina Recommendations : Tebuconazole 50% + Trifloxystrobin 25% WG is a systemic fungicide used for the control of sheath blight, leaf blast and neck blast, glume discoloration (dirty panicle) in rice crop; early blight in tomato crop; powdery mildew, anthracnose in mango crop; yellow rust, powdery mildew in wheat crop; powdery mildew in grape crop; powdery mildew, anthracnose, alternaria leaf spot in chilli crop. Cautionary statements: - 1. Not to be used for any purpose other than those subscribed on label & leaflet. 2. Destroy the container after use as directed on leaflet. 3. Avoid to mix with water bodies as product is toxic to fish and honeybees.

Aegis®

SANTIVO

Tebuconazole 50% + Trifloxystrobin 25% WG

Systemic Fungicide

A product of Aegis®

Keep out of reach of children

CAUTION

Manufactured & Marketed by

Aegis®

Aegis Agro Chemicals India Pvt. Ltd.

Factory : # 237, Main Road, Mallapally (V), Nandigaon (M), Sangareddy Dist. - 502295.

Regd. Office : Plot No. 152, MIGH Colony, Hydernagar, Medchal-Malkajgiri Dist, Hyderabad, Telangana State - 500 085.

Phone : 84080 08538 / 37

In case of any Consumer Complaints Contact:
"Manager Customer Care" Cell No.: +91 84080 08538
Email:info@aegisagro.in
Regd. Office : Plot No. 152, MIGH Colony, Hydernagar, Medchal-Malkajgiri Dist. Hyderabad. Telangana State - 500 085.

Email:info@aegisagro.in

Regn.No.

CIR-273836/2023-Tebuconazole+ Trifloxystrobin (WG) (444)-138

019/TS-IML/2016 Dt. 23-7-2016

For Agriculture use only

EMPTY PACKAGES / CONTAINERS SHOULD BE DESTROYED AFTER USE