

Triangle Bottle 125ml Size : **186 x 46.5mm** (Bottel Ref: 180ml)

GARUDA - Cyantraniliprole 7.3%w/w + Diafenthiuron 36.4%w/w SC
(Cyantraniliprole 8% w/v + Diafenthiuron 40%w/v SC) (Cyantraniliprole 80 g/L + Diafenthiuron 400 g/L-480 SC)
(Broad Spectrum Systemic Insecticide)

Chemical Composition : 1.2-bromo-2-nitropropan- 1,3-diol (30% solution in water) : 0.09 % w/w, 2 Phosphoric acid : 0.28 % w/w, 3 Potassium hydroxide (50% solution in water) : 0.28 % w/w, 4 Water : Q.S. % w/w, 5 Cyantraniliprole (a.i.) 7.3 % w/w, 6 Diafenthiuron (a.i.) : 36.4 % w/w,7 DL-propanediol-(1,2) 4.6 % w/w, 8 Polydimethylsiloxane 0.46 % w/w, 9 Polyarylphenyl ether phosphate : 1.9 % w/w, 10 Propyleneoxide/ethylene-oxide copolymer : 1.9 % w/w, 11 Naphthalenesulfonic acid, sodium salt condensed with formaldehyde 1.9 % w/w, 12 Xanthan gum 0.19 % w/w, 13 1,2-benzisothiazol-3-one (20% solution in dipropylene glycol) : 0.19 % w/w, Total : 100 % w/w. **Direction Of Use :** Measure out required quantity of the product and mix it well with a small quantity of clean water. Add the remaining quantity of the water as specified with thorough agitation for total coverage of crop with a knapsack sprayer fitted with hollow cone nozzle or any other suitable sprayer. Prepare no more spray mixture than is needed for the immediate application. Do not let the spray mixture stand overnight in the spray tank. **Crop , Dosage :** Chilli, cotton - 250ml/acre. Pre Harvesting Interval : chilli-5 days, cotton : 25 days. **Re-entry period:— Precautions :** Do not use cooking utensils for preparing the spray solution. Use stick for stirring the spray solution. Avoid contact with skin, eyes and clothing. Avoid inhalation of vapors and spray mist and wear dust mask and goggles during application. For mixing & loading and during spray application, wear appropriate personal protective equipment i.e. long-sleeved shirt and long trousers (or coveralls), impermeable footwear, eye protection and chemical resistant gloves. Do not blow spray nozzle with mouth. Do not eat, drink or smoke while handling or applying the product. Wash entire body thoroughly with soap and plenty of water and change clothes after the work is over. Wash contaminated clothing separately from other normal clothing Incase of any indisposition, stop work immediately and call doctor at the earliest. Do not spray crops under stress. Do not spray during hot hours of the day, if the foliage is wet or if rain is imminent. Do not overdose by over lapping applications or by exceeding the recommended rate. **Antidote :** No specific antidote is known. Apply symptomatic therapy, **First Aid :** Remove the affected person from the danger zone to a well-ventilated room or to fresh air In case of suspected poisoning: immediately call a physician. Show label to the physician. In case of Skin contact: Take off all contaminated clothing immediately and thoroughly wash the affected parts of the body with plenty of water. Wash contaminated clothes before re-use. If skin irritation persists, call a doctor. In case of Eye contact: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses. Immediate medical attention is required. In case of Ingestion: If swallowed, seek medical advice immediately and show this container or label to the doctor. Do not induce vomiting. In case of Inhalation: Immediately move victim to fresh air. If breathing is irregular or stopped, administer artificial respiration. Keep patient warm and at rest. Call doctor immediately. **Symptoms:** In acute toxicity studies, the formulation showed a low toxicity profile. Intoxication is only imaginable if large quantities are ingested. No Specific monitoring programs in humans have been performed. **Recommendations :** Cyantraniliprole is a novel anthranilic diamide insecticide effective against a cross spectrum of chewing and sucking pests. It is a systemic insecticide, which displays strong stomach and good contact activity. Diafenthiuron is a non systemic insecticide and acaricide, which kills larvae, nymphs and adults by contact and/or stomach action. It displays translaminar activity, which means pests located underside of leaves are controlled even if they are not directly hit by spray. One litre of this formulation contains 80-gram Cyantraniliprole and 400-gram Diafenthiuron. **CAUTIONARY STATEMENTS:-** 1. Not to be used for any purpose other than those subscribed on labels & leaflets. 2. This product is reported to be very toxic to aquatic organisms, fish and honey bee and thus the product containing molecule may be avoided near aquaculture and during active foraging period of the day.

Aegis

GARUDA VEGA

Cyantraniliprole 7.3%w/w + Diafenthiuron 36.4%w/w SC

BROAD SPECTRUM SYSTEMIC INSECTICIDE

A product of Aegis

Regn.No / रजि. नं. : **CIR-369777/2026-Cyantraniliprole + Diafenthiuron (SC) (473)-41**
Mfg.Lic.No / उत्पादक लाइसेंस. नं. : **019/TS-IML/2016 Dt. 23-7-2016**

Manufactured & Marketed by :
Aegis®
Agro Chemicals India Private Limited
AN ISO 9001 : 2015 CERTIFIED COMPANY

Aegis Agro Chemicals India Pvt. Ltd,
Factory : # 237, Main Road, Mallapally (V),
Kondapur (M), Sangareddy Dist. -502295.
फैक्टरी : २३७, मेन रोड, मल्लेपल्ली (वी), कोंडापुर (एम),
संगारेड्डी जिला, तेलंगाना-५०२२९५.
Regd. Office : Plot No. 152, NRSA Colony, Hydernagar,
Moctha Malkajgiri Dist., Hyderabad,
Telangana State - 500 085. Phone : 84980 99536 / 37
रजि. कार्यालय : प्लॉट संख्या १५२, एन.आर.एस.ए. कालोनी,
हाइदरनगर, हैदराबाद, तेलंगाना - ५०००८५.
फोन : ८४९८० ९९५३६ / ३५ | Email : info@aegisagro.in

In Case of any Consumer Complaints Contact :
"Manager Customer Care" Cell No. : +91 84980 99536
Email : info@aegisagro.in | Regd. Office Address : Plot No. 152, NRSA Colony, Hydernagar, Moctha-Malkajgiri Dist., Hyderabad, Telangana State-500085.
रिज. कार्यालय: प्लॉट न.१५२, एन.आर.एस.ए. कालोनी, हाइदरनगर, हैदराबाद, तेलंगाना - ५००००५.
फोन : ८४९८० ९९५३६ / ३५ | Email : info@aegisagro.in

For Agriculture use only. केवल कृषि उपयोग के लिए।
EMPTY PACKAGES / CONTAINERS / SHOULD BE DESTROYED AFTER USE

KEEP OUT OF THE REACH OF CHILDREN
DANGER
खतरा ಎಚ್ಚರಿಕೆ
அபாயம் ஆபாயம்

Net Weight : **125ml**
उतशुद्ध सामग्री :
Batch No :
बैच नं. :
Mfg.Date :
उत्पादन तिथि :
Exp.Date :
निम्नलिखी होने का तिथि:
Max.Retail.Price ₹ :
अधिकतम खुदरा मूल्य ₹ :
(Inclusive of all taxes)
(सभी करों सहित)
Unit Sale Price Per ml ₹ :
(यूनिट बिक्री मूल्य प्रति मिली) ₹ :

Triangle Bottle 250ml Size : **202 x 56mm**

GARUDA - Cyantraniliprole 7.3%w/w + Diafenthiuron 36.4%w/w SC
(Cyantraniliprole 8% w/v + Diafenthiuron 40%w/v SC) (Cyantraniliprole 80 g/L + Diafenthiuron 400 g/L-480 SC)
(Broad Spectrum Systemic Insecticide)

Chemical Composition : 1.2-bromo-2-nitropropan- 1,3-diol (30% solution in water) : 0.09 % w/w, 2 Phosphoric acid : 0.28 % w/w, 3 Potassium hydroxide (50% solution in water) : 0.28 % w/w, 4 Water : Q.S. % w/w, 5 Cyantraniliprole (a.i.) 7.3 % w/w, 6 Diafenthiuron (a.i.) : 36.4 % w/w,7 DL-propanediol-(1,2) 4.6 % w/w, 8 Polydimethylsiloxane 0.46 % w/w, 9 Polyarylphenyl ether phosphate : 1.9 % w/w, 10 Propyleneoxide/ethylene-oxide copolymer : 1.9 % w/w, 11 Naphthalenesulfonic acid, sodium salt condensed with formaldehyde 1.9 % w/w, 12 Xanthan gum 0.19 % w/w, 13 1,2-benzisothiazol-3-one (20% solution in dipropylene glycol) : 0.19 % w/w, Total : 100 % w/w. **Direction Of Use :** Measure out required quantity of the product and mix it well with a small quantity of clean water. Add the remaining quantity of the water as specified with thorough agitation for total coverage of crop with a knapsack sprayer fitted with hollow cone nozzle or any other suitable sprayer. Prepare no more spray mixture than is needed for the immediate application. Do not let the spray mixture stand overnight in the spray tank. **Crop , Dosage :** Chilli, cotton - 250ml/acre. Pre Harvesting Interval : chilli-5 days, cotton : 25 days. **Re-entry period:— Precautions :** Do not use cooking utensils for preparing the spray solution. Use stick for stirring the spray solution. Avoid contact with skin, eyes and clothing. Avoid inhalation of vapors and spray mist and wear dust mask and goggles during application. For mixing & loading and during spray application, wear appropriate personal protective equipment i.e. long-sleeved shirt and long trousers (or coveralls), impermeable footwear, eye protection and chemical resistant gloves. Do not blow spray nozzle with mouth. Do not eat, drink or smoke while handling or applying the product. Wash entire body thoroughly with soap and plenty of water and change clothes after the work is over. Wash contaminated clothing separately from other normal clothing Incase of any indisposition, stop work immediately and call doctor at the earliest. Do not spray crops under stress. Do not spray during hot hours of the day, if the foliage is wet or if rain is imminent. Do not overdose by over lapping applications or by exceeding the recommended rate. **Antidote :** No specific antidote is known. Apply symptomatic therapy, **First Aid :** Remove the affected person from the danger zone to a well-ventilated room or to fresh air. In case of suspected poisoning: immediately call a physician. Show label to the physician. In case of Skin contact: Take off all contaminated clothing immediately and thoroughly wash the affected parts of the body with plenty of water. Wash contaminated clothes before re-use. If skin irritation persists, call a doctor. In case of Eye contact: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses. Immediate medical attention is required. In case of Ingestion: If swallowed, seek medical advice immediately and show this container or label to the doctor. Do not induce vomiting. In case of Inhalation: Immediately move victim to fresh air. If breathing is irregular or stopped, administer artificial respiration. Keep patient warm and at rest. Call doctor immediately. **Symptoms:** In acute toxicity studies, the formulation showed a low toxicity profile. Intoxication is only imaginable if large quantities are ingested. No Specific monitoring programs in humans have been performed. **Recommendations :** Cyantraniliprole is a novel anthranilic diamide insecticide effective against a cross spectrum of chewing and sucking pests. It is a systemic insecticide, which displays strong stomach and good contact activity. Diafenthiuron is a non systemic insecticide and acaricide, which kills larvae, nymphs and adults by contact and/or stomach action. It displays translaminar activity, which means pests located underside of leaves are controlled even if they are not directly hit by spray. One litre of this formulation contains 80-gram Cyantraniliprole and 400-gram Diafenthiuron. **CAUTIONARY STATEMENTS:-** 1. Not to be used for any purpose other than those subscribed on labels & leaflets. 2. This product is reported to be very toxic to aquatic organisms, fish and honey bee and thus the product containing molecule may be avoided near aquaculture and during active foraging period of the day.

Aegis

GARUDA VEGA

Cyantraniliprole 7.3%w/w + Diafenthiuron 36.4%w/w SC

BROAD SPECTRUM SYSTEMIC INSECTICIDE

A product of Aegis

Regn.No / रजि. नं. : **CIR-369777/2026-Cyantraniliprole + Diafenthiuron (SC) (473)-41**
Mfg.Lic.No / उत्पादक लाइसेंस. नं. : **019/TS-IML/2016 Dt. 23-7-2016**

Manufactured & Marketed by :
Aegis®
Agro Chemicals India Private Limited
AN ISO 9001 : 2015 CERTIFIED COMPANY

Aegis Agro Chemicals India Pvt. Ltd,
Factory : # 237, Main Road, Mallapally (V),
Kondapur (M), Sangareddy Dist. -502295.
फैक्टरी : २३७, मेन रोड, मल्लेपल्ली (वी), कोंडापुर (एम),
संगारेड्डी जिला, तेलंगाना-५०२२९५.
Regd. Office : Plot No. 152, NRSA Colony, Hydernagar,
Moctha-Malkajgiri Dist., Hyderabad,
Telangana State - 500 085. Phone : 84980 99536 / 37
रजि. कार्यालय : प्लॉट संख्या १५२, एन.आर.एस.ए. कालोनी,
हाइदरनगर, हैदराबाद, तेलंगाना - ५०००८५.
फोन : ८४९८० ९९५३६ / ३५ | Email : info@aegisagro.in

In Case of any Consumer Complaints Contact :
"Manager Customer Care" Cell No. : +91 84980 99536
Email : info@aegisagro.in | Regd. Office Address : Plot No. 152, NRSA Colony, Hydernagar, Moctha-Malkajgiri Dist., Hyderabad, Telangana State-500085.
रिज. कार्यालय: प्लॉट न.१५२, एन.आर.एस.ए. कालोनी, हाइदरनगर, हैदराबाद, तेलंगाना - ५००००५.
फोन : ८४९८० ९९५३६ / ३५ | Email : info@aegisagro.in

For Agriculture use only. केवल कृषि उपयोग के लिए।
EMPTY PACKAGES / CONTAINERS / SHOULD BE DESTROYED AFTER USE

KEEP OUT OF THE REACH OF CHILDREN
DANGER
खतरा ಎಚ್ಚರಿಕೆ
அபாயம் ஆபாயம்

Net Weight : **250ml**
उतशुद्ध सामग्री :
Batch No :
बैच नं. :
Mfg.Date :
उत्पादन तिथि :
Exp.Date :
निम्नलिखी होने का तिथि:
Max.Retail.Price ₹ :
अधिकतम खुदरा मूल्य ₹ :
(Inclusive of all taxes)
(सभी करों सहित)
Unit Sale Price Per ml ₹ :
(यूनिट बिक्री मूल्य प्रति मिली) ₹ :